

Indian J Palliat Care. 2019 Jul-Sep; 25(3): 367–373.

doi: 10.4103/IJPC.IJPC_22_19: 10.4103/IJPC.IJPC_22_19

PMCID: PMC6659521

PMID: [31413450](#)

“Adjunctive Effects of a Short Session of Music on Pain, Low-mood and Anxiety Modulation among Cancer Patients” – A Randomized Crossover Clinical Trial

[Gunasekara Vidana Mestriige Chamath Fernando](#),^{1,2} [Lagath Udara Wanigabadu](#),³ [Buddhika Vidanagama](#),⁴ [Terancy Shyamale Perera Samaranayaka](#),^{1,2} and [Jeewandara Mudiyanseelage Kamal Chandima Jeewandara](#)^{1,2}

Address for correspondence: Dr. Gunasekara Vidana Mestriige Chamath Fernando, National Centre for Primary Care and Allergy Research, University of Sri Jayewardenepura, Nugegoda, Sri Lanka. E-mail: moc.liamg@ftemehc

Copyright : © 2019 Indian Journal of Palliative Care

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

INTRODUCTION

“Pain” is an unpleasant and a distressing feeling that affects humans physically and on occasions psychosocially. Fifty percent of all cancer patients suffer pain, while this proportion rises to 75% among those with advanced cancer.[1] Particularly, in cancer patients, the pain they suffer is closely accompanied by low mood and anxiety,[2] collectively, leading to a poor quality of life.

Cure from cancer is not a realistic goal in all instances. Relieving the patients of their symptoms seems to be a more realistic and feasible task in the majority of cases. Despite the compelling evidence to the fact that pain can be successfully alleviated in 90% of cancer patients,[3] a recent review of published literature estimated that one in every two patients were inadequately treated for cancer-related pain.[1]

The range of physical and psychological adverse effects of pain relief medications results in impairment of quality of life. This accounts for refusal of those drugs by patients and reluctance of health-care staff to prescribe or administer such medicine.[4,5,6] Fear of addiction and myriad other misconceptions borne by health-care staff and patients about the use of narcotic analgesics in terminal illnesses has also jeopardized prescription rates.[7] The development of tolerance and dependence for analgesics are known to potentially worsen pain with prolonged administration.[8] Weiss *et al.* revealed that 50% of terminal cancer patients experienced moderate-to-severe pain of whom only 30% wanted additional treatment than opioids.[6]

A range of nonpharmacological interventions reports varying degrees of success in cancer pain management, which include breathing and relaxation techniques, distraction therapies, hypnotherapy, imagery, heat or cold application, massage, pressure, vibration, and transcutaneous electrical nerve stimulation.[3,9,10] Included among these is music therapy which has been identified to have a consistent therapeutic benefit on alleviation of pain of all origins.

Music is a universal language and a multifaceted phenomenon that entails physical, emotional, cognitive, and social elements that affects people on an individual basis.[11] Music had been used for its therapeutic effects since millennia ago, of which the earliest references date back to the 5th century BC in the era of the philosophers – Plato, Socrates, and Aristotle.[12,13] They believed in the ability of music to communicate the emotional states of humans. Music has been suggested to stimulate more parts of the brain than any other human function, explaining the magnitude of belief and interest in research on music therapy.

In Canada, C. E. Dobek has assessed the effects of music on the pain response in central nervous system using functional magnetic resonance imaging. It is considered a highly accurate means of ascertaining the levels of pain experienced by people in an objective manner. It was confirmed that the emotional valence of music affects neural activity in the brainstem and spinal cord. Subjective pain ratings were significantly lower when painful stimuli were co-administered with music than without music.[11]

A study by SL Beck on the effect of the therapeutic use of music on cancer-related pain and low mood measured these dimensions on visual analog scale (VAS). The effect of music on pain demonstrated significant inter-individual variations. Seventy-five percent showed “some” response of which 47% demonstrated a “moderate” to “great” response.[14]

Aforementioned international literature suggest that music has a significant effect on pain relief. In Sri Lanka, one publication revealed how Indian classical music significantly reduced blood pressure, pulse rate, and respiratory rate in asymptomatic individuals.[15] The same group of researchers in other studies found that Indian classical music has a significant effect in relieving the symptoms and improving exercise tolerance quality of life of patients with stable angina.[16,17,18]

AIMS

Cancer patients experience pain, anxiety, and low mood, together with symptoms of related sympathetic overactivity (such as raised pulse rate, blood pressure, pupillary size, and respiratory rate).[19,20,21,22,23] Our study aimed to fill the gap in knowledge about the potential effect of culturally appropriate instrumental classical music on the alleviation of these symptoms and signs among a cohort of cancer patients in Sri Lanka.

Two related randomized controlled clinical trials conducted by Brazilian and Taiwanese investigators exclusively looked at the immediate effects of music.[24,25] They stressed the importance of determining the longevity of the effects of music on pain. Hence, our study also attempts to establish as to how long do the beneficial effects of music last on each of the subjective and objective parameters, if at all.

MATERIALS AND METHODS

Study design

This randomized, controlled, open, interventional, crossover clinical trial[26] was conducted to assess the effect of culturally familiar instrumental classical music on pain, anxiety, and low-mood modulation.

Study population

Adult resident cancer patients older than 12 years at a major tertiary care cancer institute of Sri Lanka who essentially suffered basal levels of pain (i.e., at least 1/10 on VAS) for a duration >12 weeks were included in the study.

Those who were subconscious (Glasgow Coma Scale short of 15),^[27] confused or irrational, suffering auditory impairment, not able to communicate, not suffering pain, and not on regular analgesics at all were excluded from the trial. Those whose discharge was planned within 48 h of commencement of the study and patients of surgical wards were also excluded.

Of the included patients, those who suffered “incident pain” due to interventions, activity, or mobilization, those who received additional potent analgesics or opioids to manage breakthrough pain, whose analgesic regimens altered during the course of the 48 h concerned and whose mental status deteriorated unexpectedly were planned to be removed from further assessment although the fact that such attrition did not occur.

Randomization and sample size calculation

Calculated sample size was 21 in relation to the primary outcome measures (minimum value = 0, maximum value = 10, minimally significant difference = 2, standard deviation [SD] = $10-0/4 = 2.5$ and expected difference/SD = $2/2.5 = 0.8$ for a confidence interval (CI) of 95%).^[28] We included 24 (21 + 3) patients in the study by simple randomization of which 50% were women.

The researchers took steps to identify all the patients meeting the inclusion criteria in the oncology units who were given a unique identification code. Of the total of 68 patients in the three oncology units on the initial day, 29 patients were deemed ineligible. All the remaining 39 consented participation. Twenty-four patients were selected out of the 39 eligible patients as illustrated in [Chart 1](#) through computer-assisted simple randomization.

Interventions

There were two arms concerning this study carried out in May 2017. In this open trial, the same group of patients were subjected initially to “control” and subsequently to “intervention” phase, thus conforming to the crossover design.

Duration of assessment

A pilot study on this population involving eight patients revealed that the duration of the effect of music on each measured parameter varied and none persisted beyond 12 h. Hence, it was decided to conduct the study to assess the said

parameters periodically over 24 h following exposure to music in an aim to assess the existence of an effect and if so the duration of effect for each parameter.

Depending on the duration of effect on each parameter and the symptomatology of individual patients, the frequency of listening to music was expected to be justified in the future practice.

Measurements

The primary (subjective) parameters such as pain, anxiety and grief were measured on VAS which is an accepted tool with cross-cultural validity[29] augmented with WongBaker Faces Pain Scale.[30] The addition of the latter scale improves the comprehension by individuals with marginal cognitive and mathematical abilities.[31] The VAS is primarily meant to assess the intensity of pain.[32] Nevertheless, there is a wealth of evidence now, for its use in the assessment of the degree of anxiety and low mood for which the tool has been validated.[33]

Since it was required to assess the cohort of patients 4 hourly over 48 h, the assessment tool had to be simple and least disruptive to the patients. Therefore, we did not use exhaustive questionnaires such as Brief Pain Inventory,[34] Hamilton Anxiety Rating Scale,[35] or Beck's Depression Inventory.[36]

The values for objective/surrogate measures, i.e., palpated radial pulse rate, systolic and diastolic brachial arterial blood pressure read by mercury sphygmomanometry, respiratory rate measured visually, and pupillary size observed visually under moderate daylight were also recorded.

Choice of music

International researchers employed culturally familiar musical compositions for therapy in respective regions for which T. Sathish Kumar in India utilizing Carnatic music and L. Rafer in the USA utilizing Jazz music are examples.[37,38] The music used in our study was composed by two esthetically qualified Sri Lankan composers with Western, South Indian, and Persian influence. This selection was made to ensure the cultural familiarity of the musical styles to the predominant ethnic groups – Sinhalese (74.5%), Tamil (11.15%), and Moor (9.4%) of Sri Lanka. Lyrics were not used in the composition to avoid the potential dimension of “literal meaning” acting as a confounder.

Phases

Phase A

On the 1st day, basal scores for the parameters were recorded at 08:00 AM. From 08:00 to 08:30 AM patients continued to lie in bed with their eyes gently blindfolded while they wore earplugs. After 08:30 AM, they were instructed to engage in their daily routines. The same parameters as above were recorded immediately after the rest (08:30 AM) and 4 hourly thereon.

Phase B

The same group of patients moved onto Phase B the following day, where the parameters were documented at 08:00 AM. From 08:00 to 08:30 h, patients lay in bed with their eyes gently blindfolded while they listened to a 28-min long music clip through an MP3 player with earphones that limited external noise. The volume was adjusted to a level indicated as comfortable by the patient. The parameters as above were recorded at the same time points as the previous day.[\[31,32,33\]](#)

There was no attrition (fallouts) of study participants. Once 100% of the sample had satisfactorily completed 48 h, further recruitments were deemed unnecessary.

Statistical analysis

Demographic data of the participants were described in terms of percentages. Paired sample *t*-test[\[39\]](#) was used to compare the chronologically corresponding values (16:00 h with 16:00 hrs^{+1Day}) read for individual parameters on the 2 consecutive days. This statistical procedure was intended to determine whether or not there was a mean difference in the values between Phase A (control) and Phase B (intervention). Thereof, it was aimed to assess whether the data provided evidence as for superiority of administration of classical music to conventional therapy alone in terms of symptomatic relief subjectively and the clinical parameters measured objectively. Further, the duration up to which each effect lasted (significantly) was also ascertained.

Ancillary analysis was aimed at assessing the presence of a significant difference in the measured effects, based on the patients' age, sex, and ethnicity, whether or not enjoyed listening to music and level of baseline pain.

RESULTS

None of the patients discontinued the study owing to administrative reasons, adverse effects, lack of therapeutic efficacy, or their autonomous choice. Once 100% of the sample had satisfactorily completed 48 h, further recruitments were deemed unnecessary.

Patient demographics and baseline characteristics are illustrated in [Table 1](#). The patients represented equal gender compositions, significant range in age, and all majority Sri Lankan races. Patients who suffered different levels of pain, receiving all levels of analgesics, patients who regularly enjoyed listening to music, those who did not, those averse toward pain killers and those who were not were also included. The leading types of malignancies were gastrointestinal, respiratory, gynecological and endocrine. A third had advanced stage of cancer with distant spread.

Primary end points

Analysis revealed a significant alleviation of all subjective symptoms immediately following administration of music. The levels of pain and anxiety remained significantly diminished until the 4th h reading point ($P = 0.007$, 95% CI: 3.716 to -1.534 and $P = 0.0022$, 95% CI: -2.465 to -0.6184 , respectively). The effect on depressed mood was significant until the 12th h post-administration ($P = 0.007$, 95% CI: -2.194 to -0.3896) and was absent at the 24th h assessment point.

Surrogate end points

The modulation of objective parameters was estimated as secondary end points. The pupillary size was significant up to the 12th-h measurement ($P = 0.0166$, 95% CI: -0.6752 to -0.0748) and the respiratory rate was reduced significantly for 8 h following listening to music ($P = 0.0098$, 95% CI: -0.0645 to -1.981). The fluctuations observed in hemodynamic parameters (pulse rate and blood pressure) were statistically insignificant [[Figure 1](#)].

Ancillary analysis

In the ancillary analysis, it was revealed that the patients' age, sex, ethnic background or level of baseline pain experienced did not have a significant effect on the alleviation of pain, low mood, or anxiety with music. However, anxiolysis achieved in those who did not enjoy listening to music was significantly greater than those who did ($P = 0.016$).

DISCUSSION

The participants representing the adult cancer patients receiving resident care at this tertiary care institution significantly benefitted from listening to the culturally sensitive, tranquilizing, instrumental classical music half-an-hour a day. The stated benefit was two folds in terms of all subjective parameters (pain, low mood, and anxiety) as well as related objective parameters (pupillary size and respiratory

rate). The potential of music to touch the psychosocial elements of “total pain”[40] that the conventional drugs cannot practically reach may have a bearing on this additive effect.

Generalizability and limitations

Importantly, the session did not have any adverse effects attributed to have occurred due to music on the study population which included 4 participants (16.66%) who did not regularly listen to music. The only patient who continued to suffer 10/10 pain in spite of music therapy, having been offered the option to withdraw, opted to continue and did not report additional disturbance due to music or monitoring. Hence, music therapy can be justified to be safely applied to complement conventional symptomatic treatment of adult cancer patients through a common sound system.

As mentioned before, the musical composition was deemed culturally appropriate and tranquilizing by two qualified musicians, thus enhancing the generalizability to the Sri Lankan setting.

The investigators were careful to select equal numbers of patients from the two sexes. The aforesaid exclusions were made through consensus among the investigators that the primary outcomes (involving measurement of subjective parameters on VAS) will be an area poorly comprehended by those groups. The patients in the surgical wards were not included as the chance of occurrence of incident pain (e.g., due to wound dressings, mobilization, and procedures) was expected to be higher among them.

The patients could ideally have been assessed at all 4-hourly intervals following the control and intervention. However, it was decided not to monitor them at midnight and 04:00 AM with the intention of not disturbing their sleep. A statistically significant alleviation was noted both in low mood and pupillary size at 8 PM, and the effect had disappeared by 8 AM in the following morning. Hence, the duration of persistence of these effects which fell in between 8 PM and 8 AM could not be ascertained with precision.

Another perceived limitation was the Colombo (commercial capital of Sri Lanka)-based study setting. Nevertheless, patients from all regions of the island belonging to all educational and social strata receive care in this principle tertiary care state-operated cancer institution.

CONCLUSIONS

This study was able to demonstrate that a short session of culturally familiar, instrumental classical music is capable of alleviating pain, anxiety, and low mood in institutionalized cancer patients, complementing conventional symptomatic therapies.

There would be a scope to explore in the future studies, the degree to which symptom-relieving medications can be cut down with co-administration of music while achieving the same level of symptomatic relief. The effects of longer term periodical exposure to music and sensitivities of Sri Lankan population to different genres of music are other potential areas for future investigation.

Financial support and sponsorship

This study was financially supported by Academic Student Affairs and Publications Division, University of Sri Jayewardenepura, Sri Lanka (Grant No: ASP/01/RE/MED/2016/073).

Conflicts of interest

There are no conflicts of interest.

Acknowledgments

The authors wish to acknowledge BU Gunathilaka and AC Atapattu for their contribution in composing and accreditation of the music clip along with the principal investigator (a musician himself), Director of National Cancer Institute, Sri Lanka and relevant consultants for granting permission to carry out the trial in their units.

REFERENCES

1. Deandrea S, Montanari M, Moja L, Apolone G. Prevalence of undertreatment in cancer pain. A review of published literature. *Ann Oncol.* 2008;19:1985–91. [PMCID: PMC2733110] [PubMed: 18632721]
2. Hansdorfer-Korzon R, Chojnacka-Szawłowska G, Landowski J, Majkowiec M, Basiński K, Zdun-Ryżewska A, et al. Relationships of anxiety and depressive symptoms with pain perception in postmastectomy women. An intragroup analysis. *Arch Clin Psychiatry São Paulo.* 2016;43:74–8.
3. Nersesyan H, Slavin KV. Current approach to cancer pain management: Availability and implications of different treatment options. *Ther Clin Risk Manag.* 2007;3:381–400. [PMCID: PMC2386360] [PubMed: 18488078]

4. Von Roenn JH, Cleeland CS, Gonin R, Hatfield AK, Pandya KJ. Physician attitudes and practice in cancer pain management. A survey from the Eastern Cooperative Oncology Group. *Ann Intern Med.* 1993;119:121–6. [PubMed: 8099769]
5. Grossman SA. Undertreatment of cancer pain: Barriers and remedies. *Support Care Cancer.* 1993;1:74–8. [PubMed: 7511475]
6. Weiss SC, Emanuel LL, Fairclough DL, Emanuel EJ. Understanding the experience of pain in terminally ill patients. *Lancet.* 2001;357:1311–5. [PubMed: 11343734]
7. Marks RM, Sachar EJ. Undertreatment of medical inpatients with narcotic analgesics. *Ann Intern Med.* 1973;78:173–81. [PubMed: 4683747]
8. Fields HL. The doctor's dilemma: Opiate analgesics and chronic pain. *Neuron.* 2011;69:591–4. [PMCID: PMC3073133] [PubMed: 21338871]
9. Pujol LA, Monti DA. Managing cancer pain with nonpharmacologic and complementary therapies. *J Am Osteopath Assoc.* 2007;107:ES15–21. [PubMed: 18165373]
10. Ripamonti CI, Bandieri E, Roila F. Management of cancer pain: ESMO clinical practice guidelines. *Ann Oncol.* 2011;22(Suppl 6):vi69–77. [PubMed: 21908508]
11. Dobek CE. Effects of Music on the Pain Response in the Central Nervous System Using functional Magnetic Resonance Imaging [Thesis] 2013. [Last accessed on 2017 Jun 19]. Available from: <https://www.qspace.library.queensu.ca/handle/1974/8083> .
12. Discoveries A. Music Training Confirmed by Socrates and Plato Music Influence. [Last accessed on 2017 Jun 19]. Available from: http://www.amazingdiscoveries.org/S-deception-music_philosophers_culture_plato .
13. Psychiatry and Music. [Last accessed on 2018 Jan 11]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4040058/>
14. Beck SL. The therapeutic use of music for cancer-related pain. *Oncol Nurs Forum.* 1991;18:1327–37. [PubMed: 1762973]
15. The Effect of Music on Non-Responsive Patients in a Hospice Setting. Research Gate. [Last accessed on 2017 Jun 20]. Available from: https://www.researchgate.net/publication/254673541_The_Effect_of_Music_on_Non-Responsive_Patients_in_a_Hospice_Setting .

16. Simavli S, Gumus I, Kaygusuz I, Yildirim M, Usluogullari B, Kafali H, et al. Effect of music on labor pain relief, anxiety level and postpartum analgesic requirement: A randomized controlled clinical trial. *Gynecol Obstet Invest.* 2014;78:244–50. [PubMed: 25227477]
17. Siritunga S, Wijewardena K, Ekanayaka R, Mudunkotuwa P. Effect of music on blood pressure, pulse rate and respiratory rate of asymptomatic individuals: A randomized controlled trial. *Health (N Y)* 2013;5:59.
18. Siritunga S, Wijewardena K, Ekanayaka R, Mudunkotuwa P. Can music improve the symptoms of stable angina? A randomized controlled trial. *Health (NY)* 2013;5:1039.
19. Siritunga S, Wijewardena K, Ekanayaka R, Mudunkotuwa P. Effect of music on state of ischaemia in stable angina; a randomized controlled trial. *Int J Clin Med.* 2014;5:1173.
20. Siritunga S, Wijewardena K, Ekanayaka R, Mudunkotuwa P. Effect of music on quality of life in stable angina: A randomized controlled trial. *Int J Clin Med.* 2015;6:307.
21. Brown LF, Kroenke K, Theobald DE, Wu J, Tu W. The association of depression and anxiety with health-related quality of life in cancer patients with depression and/or pain. *Psychooncology.* 2010;19:734–41. [PMCID: PMC2888919] [PubMed: 19777535]
22. Nicolson SE, Caplan JP, Williams DE, Stern TA. Comorbid pain, depression, and anxiety: Multifaceted pathology allows for multifaceted treatment. *Harv Rev Psychiatry.* 2009;17:407–20. [PubMed: 19968455]
23. Deng G, Cassileth BR. Integrative oncology: Complementary therapies for pain, anxiety, and mood disturbance. *CA Cancer J Clin.* 2005;55:109–16. [PubMed: 15761079]
24. Jarrett ME, Burr RL, Cain KC, Hertig V, Weisman P, Heitkemper MM, et al. Anxiety and depression are related to autonomic nervous system function in women with irritable bowel syndrome. *Dig Dis Sci.* 2003;48:386–94. [PubMed: 12643620]
25. Crofford LJ. Chronic pain: Where the body meets the brain. *Trans Am Clin Climatol Assoc.* 2015;126:167–83. [PMCID: PMC4530716] [PubMed: 26330672]
26. SLCTR. [Last accessed on 2017 Oct 28]. Available from: <http://www.slctr.lk/trials/358> .

27. Glasgow Coma Scale. [Last accessed on 2017 Oct 09]. Available from: <http://www.glasgowcomascale.org/>
28. Re: st: Sample Size Calculation. [Last accessed on 2017 Oct 29]. Available from: <https://www.stata.com/statalist/archive/2005-10/msg00806.html> .
29. Carlsson AM. Assessment of chronic pain. I. Aspects of the reliability and validity of the visual analogue scale. *Pain*. 1983;16:87–101. [PubMed: 6602967]
30. Paired Sample *T*-Test – Statistics Solutions. [Last accessed on 2017 Oct 28]. Available from: <http://www.statisticssolutions.com/manova-analysis-paired-sample-t-test/>
31. Ping E. Wong-Baker Faces Scale. [cited 2019 Mar 26]. Available from: https://www.academia.edu/7329990/Wong-Baker_Faces_Scale .
32. Boonstra AM, Stewart RE, Köke AJA, Oosterwijk RFA, Swaan JL, Schreurs KMG, et al. Cut-Off Points for Mild, Moderate, and Severe Pain on the Numeric Rating Scale for Pain in Patients with Chronic Musculoskeletal Pain: Variability and Influence of Sex and Catastrophizing. *Front Psychol* [Internet] [cited 2017 Jun 21];2016 7 Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5043012/> [PMCID: PMC5043012] [PubMed: 27746750]
33. Williams VS, Morlock RJ, Feltner D. Psychometric evaluation of a visual analog scale for the assessment of anxiety. *Health Qual Life Outcomes*. 2010 Jun;8:8–57. [PMCID: PMC2904728] [PubMed: 20529361]
34. Cleeland CS, Ryan KM. Pain assessment: Global use of the brief pain inventory. *Ann Acad Med Singapore*. 1994;23:129–38. [PubMed: 8080219]
35. Hamilton M. The assessment of anxiety states by rating. [Last accessed on 2017 Jun 21];*Br J Med Psychol* [Internet] 1959 32 Available from: <https://doi.org/10.1111/j.2044-8341.1959.tb00467.x>.
36. What Exactly is the Mughal/Persian Influence on Hindustani Classical Music? Quora. [Last accessed on 2017 Oct 29]. Available from: <https://www.quora.com/What-exactly-is-the-Mughal-Persian-influence-on-Hindustani-classical-music> .
37. Kumar TS, Muthuraman M, Krishnakumar R. Effect of the raga ananda bhairavi in post operative pain relief management. *Indian J Surg*. 2014;76:363–70. [PMCID: PMC4571510] [PubMed: 26396469]

38. Rafer L, Austin F, Frey J, Mulvey C, Vaida S, Prozesky J, et al. Effects of jazz on postoperative pain and stress in patients undergoing elective hysterectomy. *Adv Mind Body Med*. 2015;29:6–11. [PubMed: 25607117]
39. Othman AR, Wilcox RR, Fradette K The New and Improved Two-Sample t Test - Keselman HJ. 2004. [Cited 2019 Mar 15]. [Internet] Available from: <https://journals.sagepub.com/doi/abs/10.1111/j.0963-7214.2004.01501008.x> . [PubMed: 14717831]
40. ChiKeong O, Duncan F. Embracing Cicely Saunders's concept of total pain. *BMJ*. 2005;331:576. [PMCID: PMC1200625] [PubMed: 16150775]